

Dr. Robert Koch Dr. Scott Applegate 4230 Pioneer Woods Drive Lincoln, NE 68506 402.488.7337

ACETAMINOPHEN DOSAGE CHART

• The proper dosage for your child is based on weight.

- Always use the measuring device that comes with the medicine, not a spoon from the kitchen.
- Do not exceed five doses in 24 hours.
- Do not use extra strength acetaminophen in children unless directed by the doctor.

leight: 60 to 71 pounds D mg dren's liquid: 2 ½ tsp (12.5 ml in cup) dren's chews or meltaways: 5 or strength chews or meltaways: 2 ½ It regular strength tabs (325mg): 1
/eight: 96+ pounds D mg dren's liquid: 4 tsp (20 ml in cup) dren's chews or meltaways: 8 or strength chews or meltaways: 4 It regular strength tabs (325mg): 2

IBUPROPHEN DOSAGE CHART

- The proper dosage for your child is based on weight.
- Always use the measuring device that comes with the medicine, not a spoon from the kitchen.
- You can repeat the dose every 6 to 8 hours. Do not exceed four doses in 24 hours.

Child's Weight: 24 to 35 pounds Dose: 100 mg Infant drops: 2 droppers OR 1 1/3 syringe (2.5 ml) Children's liquid: 1 tsp	Child's Weight: 36 to 47 pounds Dose: 150 mg Infant drops: 3 droppers OR 2 syringes (3.75ml) Children's liquid: 1 ½ tsp Children's chews: 3
Child's Weight: 48 to 59 pounds Dose: 200 mg Children's liquid: 2 tsp Children's chews: 4 Junior strength chews or caplets: 2	Child's Weight: 60 to 71 pounds Dose: 250 mg Children's liquid: 2 ½ tsp Children's chew: 5 Junior strength chews or caplets: 2 ½ Adult regular strength tabs: 1
Child's Weight: 72 to 95 pounds Dose: 300 mg Children's liquid: 3 tsp Children's chews: 6 Junior strength chews or caplets: 3 Adult regular strength tabs: 1 ½	Child's Weight: 96+ pounds Dose: 400 mg Children's liquid: 4 tsp Children's chews: 8 Junior strength chews or caplets: 4 Adult regular strength tabs: 2

A fever is defined as any temperature over 100.4° Fahrenheit (38° Celsius).