

A fever is defined as any temperature over 100.4^o Fahrenheit (38^o Celsius).

ACETAMINOPHEN DOSAGE CHART

- The proper dosage for your child is based on weight.
- Do not give to a baby under 3 months old without your doctor's approval.
- Always use the measuring device that comes with the medicine, not a spoon from the kitchen.
- Don't confuse infant drops with children's liquid. Infant drops come with a dropper and are much more concentrated. The infant drops have been discontinued but may still be in your home.
- Do not exceed five doses in 24 hours.

Child's Weight: 6 to 11 pounds Dose: 40 mg Infant liquid: 1.25 ml in syringe	Child's Weight: 12 to 17 pounds Dose: 80 mg Infant liquid: 2.5 ml in syringe Children's liquid: ½ tsp (2.5 ml in cup)
Child's Weight: 18 to 23 pounds Dose: 120 mg Infant liquid: 3.75 ml in syringe Children's liquid: ¾ tsp (3.75 ml in cup) Children's chews or meltaways: 1 ½	Child's Weight: 24 to 35 pounds Dose: 160 mg Infant liquid: 5 ml in syringe Children's liquid: 1 tsp (5 ml in cup) Children's chews or meltaways: 2 Jr. Strength chews or meltaways: 1

Your child weighs _____ pounds (_____ %tile) and is _____ tall (_____ %tile).